

Pre-Treatment*Instructions for all Procedures:

Anna Pezina (sole trader) trading as "Beaushade"
ABN 81440818275

1. Do not take any anti-inflammatory remedies such as aspirin, paracetamol etc.
2. Please do not drink alcohol 24 hours prior to the treatment.
3. Consult your doctor for any specific concerns such as recent surgery or current medications, particularly blood thinners.
4. Contact lenses should not be worn during an eyeliner procedure and 2 days after it.
5. Please do not shape or wax your brows before the procedure. Your technician will shape brows during the procedure.
6. Botox, AHA products and retinoids should be avoided for 2 weeks prior to the procedure.
7. Exfoliating treatments such as microdermabrasion should not be performed within 2 weeks prior to procedure.
8. Chemical and laser peels should be avoided no less than 6 weeks prior to procedure.
9. Patients prone to cold sores/fever blisters are advised to take vitamins prior to treatment and an anti-viral medicine after treatment.
10. Hormone therapies can affect pigmentation and/or cause sensitivity.
11. Please be sure that you have no period for eyeliner and lips treatment, it can cause sensitivity.
12. Please be sure that you are not pregnant.
13. If you are breast feeding mum please be sure that you feed baby for at least 6 month without any instability.
14. Check the list of contraindications:
 - any virus and bacterial diseases in an active stage, immune system disorders: autoimmune diseases, immunodeficiency, including HIV, herpes in exacerbation stages, virus/allergic conjunctivitis, psoriasis, neurodermatitis, allergic diseases, cancer diseases,
 - allergy to topical anesthesia components such as Lidocaine, Prilocaine, Benzocaine, Tetracaine and Adrenaline (Epinephrine)
 - allergy to metals such as Iron, Titanium, Nickel which can be in composition of some pigments.
 - dermal diseases, atypical changes of integuments, including benign and malignant tumors, pigmentation abnormalities, including erratic ones,
 - diabetes mellitus, any serious illness of internals a renal, liver failure,
 - predilection of a skin to form keloids,
 - hemophilia and other abnormalities of coagulability of a blood, high blood pressure,
 - alcoholic or drug intoxication, epilepsy or mental disorders,
 - hormonal diseases, pregnancy and breastfeeding period (especially in the beginning),
 - therapy period of antibiotics, hormonal drugs, drugs influencing change of coagulability of a blood.

If the you are under one of these states it is recommended to cancel the procedure or proceed with written consent of the doctor.